

WHAT IS CLAIMED IS:

1. An isolated polypeptide comprising an amino acid sequence selected from the group consisting of:
 - (a) a mature form of an amino acid sequence selected from the group consisting of SEQ ID NOS:2, 4, 6, 8, 13, 15, 17, 19, 21, 23, and 25;
 - (b) a variant of a mature form of an amino acid sequence selected from the group consisting of SEQ ID NOS:2, 4, 6, 8, 13, 15, 17, 19, 21, 23, and 25, wherein one or more amino acid residues in said variant differs from the amino acid sequence of said mature form, provided that said variant differs in no more than 15% of the amino acid residues from the amino acid sequence of said mature form;
 - (c) an amino acid sequence selected from the group consisting of SEQ ID NOS:2, 4, 6, 8, 13, 15, 17, 19, 21, 23, and 25; and
 - (d) a variant of an amino acid sequence selected from the group consisting of SEQ ID NOS:2, 4, 6, 8, 13, 15, 17, 19, 21, 23, and 25, wherein one or more amino acid residues in said variant differs from the amino acid sequence of said mature form, provided that said variant differs in no more than 15% of amino acid residues from said amino acid sequence.
- 2 The polypeptide of claim 1, wherein said polypeptide comprises the amino acid sequence of a naturally-occurring allelic variant of an amino acid sequence selected from the group consisting of SEQ ID NOS:2, 4, 6, 8, 13, 15, 17, 19, 21, 23, and 25.
3. The polypeptide of claim 2, wherein said allelic variant comprises an amino acid sequence that is the translation of a nucleic acid sequence differing by a single nucleotide from a nucleic acid sequence selected from the group consisting of SEQ ID NOS:1, 3, 5, 7, 9, 10, 11, 12, 14, 16, 18, 20, 22, and 24.
4. The polypeptide of claim 1, wherein the amino acid sequence of said variant comprises a conservative amino acid substitution.

5. An isolated nucleic acid molecule comprising a nucleic acid sequence encoding a polypeptide comprising an amino acid sequence selected from the group consisting of:

- (a) a mature form of an amino acid sequence selected from the group consisting of SEQ ID NOS:2, 4, 6, 8, 13, 15, 17, 19, 21, 23, and 25;
- (b) a variant of a mature form of an amino acid sequence selected from the group consisting of SEQ ID NOS:2, 4, 6, 8, 13, 15, 17, 19, 21, 23, and 25, wherein one or more amino acid residues in said variant differs from the amino acid sequence of said mature form, provided that said variant differs in no more than 15% of the amino acid residues from the amino acid sequence of said mature form;
- (c) an amino acid sequence selected from the group consisting of SEQ ID NOS:2, 4, 6, 8, 13, 15, 17, 19, 21, 23, and 25;
- (d) a variant of an amino acid sequence selected from the group consisting of SEQ ID NOS:2, 4, 6, 8, 13, 15, 17, 19, 21, 23, and 25, wherein one or more amino acid residues in said variant differs from the amino acid sequence of said mature form, provided that said variant differs in no more than 15% of amino acid residues from said amino acid sequence;
- (e) a nucleic acid fragment encoding at least a portion of a polypeptide comprising an amino acid sequence chosen from the group consisting of SEQ ID NOS:2, 4, 6, 8, 13, 15, 17, 19, 21, 23, and 25, or a variant of said polypeptide, wherein one or more amino acid residues in said variant differs from the amino acid sequence of said mature form, provided that said variant differs in no more than 15% of amino acid residues from said amino acid sequence; and
- (f) a nucleic acid molecule comprising the complement of (a), (b), (c), (d) or (e).

6. The nucleic acid molecule of claim 5, wherein the nucleic acid molecule comprises the nucleotide sequence of a naturally-occurring allelic nucleic acid variant.

7. The nucleic acid molecule of claim 5, wherein the nucleic acid molecule encodes a polypeptide comprising the amino acid sequence of a naturally-occurring polypeptide variant.

8. The nucleic acid molecule of claim 5, wherein the nucleic acid molecule differs by a single nucleotide from a nucleic acid sequence selected from the group consisting of SEQ ID NOS:1, 3, 5, 7, 9, 10, 11, 12, 14, 16, 18, 20, 22, and 24.

9. The nucleic acid molecule of claim 5, wherein said nucleic acid molecule comprises a nucleotide sequence selected from the group consisting of:
 - (a) a nucleotide sequence selected from the group consisting of SEQ ID NOS:1, 3, 5, 7, 9, 10, 11, 12, 14, 16, 18, 20, 22, and 24;
 - (b) a nucleotide sequence differing by one or more nucleotides from a nucleotide sequence selected from the group consisting of SEQ ID NOS:1, 3, 5, 7, 9, 10, 11, 12, 14, 16, 18, 20, 22, and 24, provided that no more than 20% of the nucleotides differ from said nucleotide sequence;
 - (c) a nucleic acid fragment of (a); and
 - (d) a nucleic acid fragment of (b).
10. The nucleic acid molecule of claim 5, wherein said nucleic acid molecule hybridizes under stringent conditions to a nucleotide sequence chosen from the group consisting of SEQ ID NOS:1, 3, 5, 7, 9, 10, 11, 12, 14, 16, 18, 20, 22, and 24, or a complement of said nucleotide sequence.
11. The nucleic acid molecule of claim 5, wherein the nucleic acid molecule comprises a nucleotide sequence selected from the group consisting of:
 - (a) a first nucleotide sequence comprising a coding sequence differing by one or more nucleotide sequences from a coding sequence encoding said amino acid sequence, provided that no more than 20% of the nucleotides in the coding sequence in said first nucleotide sequence differ from said coding sequence;
 - (b) an isolated second polynucleotide that is a complement of the first polynucleotide; and
 - (c) a nucleic acid fragment of (a) or (b).
12. A vector comprising the nucleic acid molecule of claim 11.
13. The vector of claim 12, further comprising a promoter operably-linked to said nucleic acid molecule.
14. A cell comprising the vector of claim 12.
15. An antibody that binds immunospecifically to the polypeptide of claim 1.

16. The antibody of claim 15, wherein said antibody is a monoclonal antibody.
17. The antibody of claim 15, wherein the antibody is a humanized antibody.
18. A method for determining the presence or amount of the polypeptide of claim 1 in a sample, the method comprising:
- (a) providing the sample;
 - (b) contacting the sample with an antibody that binds immunospecifically to the polypeptide; and
 - (c) determining the presence or amount of antibody bound to said polypeptide, thereby determining the presence or amount of polypeptide in said sample.
19. A method for determining the presence or amount of the nucleic acid molecule of claim 5 in a sample, the method comprising:
- (a) providing the sample;
 - (b) contacting the sample with a probe that binds to said nucleic acid molecule; and
 - (c) determining the presence or amount of the probe bound to said nucleic acid molecule, thereby determining the presence or amount of the nucleic acid molecule in said sample.
20. The method of claim 19 wherein presence or amount of the nucleic acid molecule is used as a marker for cell or tissue type.
21. The method of claim 20 wherein the cell or tissue type is cancerous.
22. A method of identifying an agent that binds to a polypeptide of claim 1, the method comprising:
- (a) contacting said polypeptide with said agent; and
 - (b) determining whether said agent binds to said polypeptide.
23. The method of claim 22 wherein the agent is a cellular receptor or a downstream effector.
24. A method for identifying an agent that modulates the expression or activity of the polypeptide of claim 1, the method comprising:

- (a) providing a cell expressing said polypeptide;
(b) contacting the cell with said agent, and
(c) determining whether the agent modulates expression or activity of said polypeptide,
whereby an alteration in expression or activity of said peptide indicates said agent modulates expression or activity of said polypeptide.

25. A method for modulating the activity of the polypeptide of claim 1, the method comprising contacting a cell sample expressing the polypeptide of said claim with a compound that binds to said polypeptide in an amount sufficient to modulate the activity of the polypeptide.

26. A method of treating or preventing a FCTR_X-associated disorder, said method comprising administering to a subject in which such treatment or prevention is desired the polypeptide of claim 1 in an amount sufficient to treat or prevent said FCTR_X-associated disorder in said subject.

27. The method of claim 26 wherein the disorder is a neurodegenerative disorder.

28. The method of claim 26 wherein the disorder is related to cell signal processing and metabolic pathway modulation.

29. The method of claim 26, wherein said subject is a human.

30. A method of treating or preventing a FCTR_X-associated disorder, said method comprising administering to a subject in which such treatment or prevention is desired the nucleic acid of claim 5 in an amount sufficient to treat or prevent said FCTR_X-associated disorder in said subject.

31. The method of claim 30 wherein the disorder is a neurodegenerative disorder.

32. The method of claim 30 wherein the disorder is related to cell signal processing and metabolic pathway modulation.

33. The method of claim 30, wherein said subject is a human.

34. A method of treating or preventing a FCTR_X-associated disorder, said method comprising administering to a subject in which such treatment or prevention is desired the antibody of claim 15 in an amount sufficient to treat or prevent said FCTR_X-associated disorder in said subject

35. The method of claim 34 wherein the disorder is selected from the group consisting of Also within the scope of the invention is the use of a Therapeutic in the manufacture of a medicament for treating or preventing disorders or syndromes including, *e.g.*, Colorectal cancer, adenomatous polyposis coli, myelogenous leukemia, congenital neonatal alloimmune thrombocytopenia, multiple human solid malignancies, malignant ovarian tumours particularly at the interface between epithelia and stroma, malignant brain tumors, mammary tumors, human gliomas, astrocytomas, mixed glioma/astrocytomas, renal cells carcinoma, breast adenocarcinoma, ovarian cancer, melanomas, renal cell carcinoma, clear cell and granular cell carcinomas, autocrine/paracrine stimulation of tumor cell proliferation, autocrine/paracrine stimulation of tumor cell survival and tumor cell resistance to cytotoxic therapy, paranechmal and basement membrane invasion and motility of tumor cells thereby contributing to metastasis, tumor-mediated immunosuppression of T-cell mediated immune effector cells and pathways resulting in tumor escape from immune surveillance, neurological disorders, neurodegenerative disorders, nerve trauma, familial myelodysplastic syndrome, Charcot-Marie-Tooth neuropathy, demyelinating Gardner syndrome, familial myelodysplastic syndrome; mental health conditions, immunological disorders, allergy and infection, asthma, bronchial asthma, Avellino type eosinophilia, lung diseases, reproductive disorders, male infertility, female reproductive system disorders, male and female reproductive diseases, hemangioma, deafness, glycoprotein Ia deficiency, desmoid disease, turcot syndrome, liver cirrhosis, hepatitis C, gastric disorders, pancreatic diseases like diabetes, Schistosoma mansoni infection, Spinocerebellar ataxia, Plasmodium falciparum parasitemia, Corneal dystrophy - Groenouw type I, Corneal dystrophy - lattice type I, and Reis-Bucklers corneal dystrophy.

36. The method of claim 34 wherein the disorder is related to cell signal processing and metabolic pathway modulation.

37. The method of claim 34, wherein the subject is a human.

38. A pharmaceutical composition comprising the polypeptide of claim 1 and a pharmaceutically-acceptable carrier.

39. A pharmaceutical composition comprising the nucleic acid molecule of claim 5 and a pharmaceutically-acceptable carrier.

40. A pharmaceutical composition comprising the antibody of claim 15 and a pharmaceutically-acceptable carrier.

41. A kit comprising in one or more containers, the pharmaceutical composition of claim 38.

42. A kit comprising in one or more containers, the pharmaceutical composition of claim 39.

43. A kit comprising in one or more containers, the pharmaceutical composition of claim 40.

44. A method for determining the presence of or predisposition to a disease associated with altered levels of the polypeptide of claim 1 in a first mammalian subject, the method comprising:

- (a) measuring the level of expression of the polypeptide in a sample from the first mammalian subject; and
 - (b) comparing the amount of said polypeptide in the sample of step (a) to the amount of the polypeptide present in a control sample from a second mammalian subject known not to have, or not to be predisposed to, said disease;
- wherein an alteration in the expression level of the polypeptide in the first subject as compared to the control sample indicates the presence of or predisposition to said disease.

45. The method of claim 44 wherein the predisposition is to cancers.

46. A method for determining the presence of or predisposition to a disease associated with altered levels of the nucleic acid molecule of claim 5 in a first mammalian subject, the method comprising:

- (a) measuring the amount of the nucleic acid in a sample from the first mammalian subject; and
- (b) comparing the amount of said nucleic acid in the sample of step (a) to the amount of the nucleic acid present in a control sample from a second mammalian subject known not to have or not be predisposed to, the disease;

wherein an alteration in the level of the nucleic acid in the first subject as compared to the control sample indicates the presence of or predisposition to the disease.

47. The method of claim 46 wherein the predisposition is to cancers.

48. A method of treating a pathological state in a mammal, the method comprising administering to the mammal a polypeptide in an amount that is sufficient to alleviate the pathological state, wherein the polypeptide is a polypeptide having an amino acid sequence at least 95% identical to a polypeptide comprising an amino acid sequence of at least one of SEQ ID NOS:2, 4, 6, 8, 13, 15, 17, 19, 21, 23, and 25, or a biologically active fragment thereof.

49. A method of treating a pathological state in a mammal, the method comprising administering to the mammal the antibody of claim 15 in an amount sufficient to alleviate the pathological state.